LESSON 11.3 Treating Alcohol Use Disorders Note-Taking Guide

Alcohol Use Disorder

1. Alcohol use disorder (AUD) is a(n)	condition when a person's	
drinking causes	or	
2. AUD ranges from mild to moderate to		

Possible Causes of Alcohol Use Disorder in Teens

3. Describe three of the most common reasons for teen alcohol use disorders by completing the information in the diagram.

Genetics	factors	Social factors
If a teen has a close with an alcohol problem, they may be to alcohol being pleasurable regardless of the side effects.	Teens with and depression may be predisposed to an alcohol use disorder.	Having who drink regularly; portrayals of alchol being used to have fun; and the of peers, family members, and other role models.

Getting Help for an Alcohol Use Disorder

	Indications you should seek help for an alcohol use disorder include having problems in or with:

5.	If you have a friend or a family member who has an alcohol use disorder,	they need
6.	Talk with the person with the problem when they are	_·
۱lc	cohol Treatment Programs	
7.	An alcohol treatment program is where people go to getalcohol use disorder.	help for ar

8. The type of treatment is based on the ______ of the alcohol use disorder

Outpatient Treatment

and _____ needs.

9. Complete the diagram below to explain what outpatient treatment includes.

	Students can go to
	and live at
	They have to attend
	which is set by a
Outpatient treatment	abuse counselor or
	health therapist.
	May include individual counseling,
	therapy, mutual
	groups, or student
	programs.

10. Identify what a student assistance program does by completing the diagram.

Substance abuse counselors	Brought into the	Meet with students or as a		
Student assistance	Provide early, tobacco, and	The programs are for		
programs	drug and intervention services	families.		
Student assistance programs	Assistance in getting students back into	Provide informational about		
		how to assist students in need		
Medications 11. Two of the most comi	Medications11. Two of the most commonly used medications given to stop and reduce drinking follow.			
		•		
	nuse) makes the person feel up when they drink.	or causes them to		
2. Naltrexone (Revia	a) reduces a person's	to drink.		
Residential Treatme	ent Centers			
12.				
a. A residential treatment center requires the person to live time for the length of their treatment.				
b. The length of trea	atment depends on the person's	in the program.		
c. Treatment may in as other therapeu	clude and utic and act	d group counseling sessions as well tivities.		
Family Involvemen	nt in Residential Treatment	t Programs		
	in support groups for families of pe			
1				
2				